


A comprehensive assessment of physical, psychological, and social or role functions for mentally competent ambulatory patients

Alan M. Jette, and Paul D. Cleary,
- Functional Disability Assessment, PTJournal

The FSQ was designed as a clinical tool to screen for functional disability and to monitor change in function. Automated reports: improve collaboration with psychiatric and other professional staff, assure quality of service for appropriate level of care, and facilitate stabilization.



Patients or caregivers identify behavioral health needs before the visit.

Real-time Reports

The FSQ real-time report integrates seamlessly into busy practices; It is as easy as ordering a lab test.

... Highest reliability is achieved in the basic ADL, intermediate ADL and mental health scales, says Jette et.al, "Functional Disability Assessment," p. 1856. Reliability is above .60.

Consumers answer screening questions and then self-administer tests on the non-technical interface in minutes.

BHC's and Providers immediately receive numerical scores that measure severity of a patient's symptoms.

Graphs signal composite scores

Tools consistently probe in detail


Test items show consumer's responses

Instant Excel export for grant reporting

Real-time trended-reports monitor consumer's progress

FSQ Case-Level Report
Functional Status Questionnaire

Dupont Linhoff
Patient Assessments



Name: 12345678
ID: 5/12/2009 12:12:38 PM
Patient Gender: Female Age: 43
Caregiver Ring: 5/23/2008 7/05/2008 5/12/2009

Parameter	Warning Zone	Good
Basic activities of daily living	0 - 87	88 - 100
Intermediate activities of daily living	0 - 77	78 - 100
Mental health	0 - 70	71 - 100
Work performance	0 - 78	79 - 100
Social activities	0 - 78	79 - 100
Quality of interactions	0 - 69	70 - 100

Physical Function (Activities of Daily Living, or ADL)

Basic ADL: During the past month have you had difficulty with

- (1) Taking care of yourself, that is, eating, dressing or bathing?
 - (2) Moving in or out of a bed or chair?
 - (3) Walking indoors, such as around your home?
- Response/Points: usually did with no difficulty 4, some difficulty 3, much difficulty 2, usually did not do because of healthy 1, usually did not do for other reason 0. Assume that "usually did not do for other reasons" is not a "valid" response, so that the ((maximum response score) - (minimum response score)) = 3

Some difficulty
Much difficulty
Usually did not

Intermediate ADL: During the past month have you had difficulty with


- (1) Walking several blocks?
 - (2) Walking one block or climbing one flight of stairs?
 - (3) Doing work around the house, such as cleaning, light yard work or home maintenance?
 - (4) Doing errands such as grocery shopping?
 - (5) Driving a car or using public transportation?
 - (6) Doing vigorous activities such as running, lifting heavy objects or participating in strenuous sports?
- Response/Points: usually did with no difficulty 4, some difficulty 3, much difficulty 2, usually did not do because of healthy 1, usually did not do for other reason 0. Assume that "usually did not do for other reasons" is not a "valid" response, so that the ((maximum response score) - (minimum response score)) = 3

Usually did with no difficulty
Some difficulty
Much difficulty
Usually did not
Some difficulty
Usually did not do for other reason

Psychological Function (Mental Health) During the past month,

- (1) Have you been a very nervous person?
- (2) Have you felt calm and peaceful?
- (3) Have you felt downhearted and blue?

Most of the time
A good bit of the time
Some of the time
A little of the time



Physical Function (Activities of Daily Living, or ADL)

Basic ADL: During the past month have you had difficulty with

- (1) Taking care of yourself, that is, eating, dressing or bathing?
 - (2) Moving in or out of a bed or chair?
 - (3) Walking indoors, such as around your home?
- Response/Points: usually did with no difficulty 4, some difficulty 3, much difficulty 2, usually did not do because of healthy 1, usually did not do for other reason 0. Assume that "usually did not do for other reasons" is not a "valid" response, so that the ((maximum response score) - (minimum response score)) = 3

Some difficulty
Much difficulty

THE FAQ SCALES ARE:

- Basic activities of daily living,
- Intermediate activities of daily living,
- Mental health,
- Work performance,
- Social activity and,
- Quality of interaction.

- Intermediate ADL:** During the past month have you had difficulty with
- (1) Walking several blocks?
 - (2) Walking one block or climbing one flight of stairs?
 - (3) Doing work around the house, such as cleaning, light yard work or home maintenance?
 - (4) Doing errands such as grocery shopping?
 - (5) Driving a car or using public transportation?
 - (6) Doing vigorous activities such as running, lifting heavy objects or participating in strenuous sports?
- Response/Points: usually did with no difficulty 4, some difficulty 3, much difficulty 2, usually did not do because of healthy 1, usually did not do for other reason 0. Assume that "usually did not do for other reasons" is not a "valid" response, so that the ((maximum response score) - (minimum response score)) = 3

Usually did with no difficulty
Some difficulty
Much difficulty
Usually did not
Some difficulty
Usually did not do for other reason

Psychological Function

- (1) Have you been a very nervous person?
 - (2) Have you felt calm and peaceful?
 - (3) Have you felt downhearted and blue?
 - (4) Were you a happy person?
 - (5) Do you feel so down?
- Response to 1, 3 and 5 Points: all of the time 1, most of the time 2, a good bit of the time 3, some of the time 4, a little of the time 5, none of the time 6
Response to 2 and 4 Points: all of the time 6, most of the time 5, a good bit of the time 4, some of the time 3, a little of the time 2, none of the time 1
Response to 1, 3 and 5 Points: all of the time 1, most of the time 2, some of the time 3, none of the time 4
Response to 2, 4 and 6 Points: all of the time 4, most of the time 3, some of the time 2, none of the time 1
* (Maximum response score) - (minimum response score) = 3. The subgroups of questions are scored in reverse. 0% is poor and 100% is good, reasons 0

Most of the Time
Some of the Time
All of the Time
Most of the Time

Procedure

1. Consumers self administer test in the hand-held tablet
2. Web-communication base sends data HIPAA compliant
3. BHC's receive the trended-FSQ lab report
4. Consumer data flows into the electronic record

Excel Export

5. Excel exports are instantly available for off-site research
6. Report FSQ scores that drop or raise by 3 or 5



Dupont Linhoff Patient Assessments

5827 South Reed Way, Suite 1528, Littleton, Colorado 80123, Joe@Linhoff.Com, WWW.DLPA.Biz